



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

EXPRESSIONS FOR CERTAIN ACCELERATIONS OF A PARTICLE.

By GEORGE H. CRESSE, U. S. Naval Academy.

The object of this paper is to present a method of obtaining expressions for the magnitudes of certain accelerations of a particle in terms of its speed and from our knowledge of its path. Let the path be plane and considered only at non-singular points, and let its equation be $f(x, y) = 0$. By differentiation with respect to time, we see that the motion of the particle satisfies the equation,

$$(1) \quad f_x'x' + f_y'y' = 0$$

where x' , y' are time-derivatives. By differentiating (1) with respect to time and dividing by $\sqrt{f_x'^2 + f_y'^2}$, we have,

$$(2) \quad \frac{f_x''x'^2 + 2f_{xy}'x'y' + f_y''y'^2}{\sqrt{f_x'^2 + f_y'^2}} + \frac{f_x'x'' + f_y'y''}{\sqrt{f_x'^2 + f_y'^2}} = 0.$$

where x'' , y'' are second time-derivatives. In the light of (1), we may put

$$\frac{f_x'x'' + f_y'y''}{\sqrt{f_x'^2 + f_y'^2}} = \frac{-y'x'' + x'y''}{\sqrt{x'^2 + y'^2}} = \alpha \cos \theta,$$

where α is the magnitude of the acceleration, and θ is the angle between the normal to the curve and the direction of acceleration. Hence for (2), we may write,

$$(3) \quad \frac{f_x''x'^2 + 2f_{xy}'x'y' + f_y''y'^2}{\sqrt{f_x'^2 + f_y'^2}} = -\alpha \cos \theta.$$

I. For example, let $f(x, y) = x^2 + y^2 - r^2$, and let the acceleration be directed toward the center of the circle. (3) now becomes the familiar formula:

$$(4) \quad \frac{x'^2 + y'^2}{\sqrt{x^2 + y^2}} = \alpha, \quad \text{or} \quad \alpha = \frac{v^2}{r}.$$

II. For another example, let $f(x, y) = b^2x^2 + a^2y^2 - a^2b^2$, and let the acceleration be directed toward one focus F of the ellipse. For this example, (3) is

$$(5) \quad \frac{b^2x'^2 + a^2y'^2}{\sqrt{b^4x^2 + a^4y^2}} = -\alpha \cos \theta.$$

If r and r' be the focal distances from F and F' respectively to the particle at P , and p and p' be perpendiculars drawn from the foci to the tangent which has

P for its point of tangency, elementary properties of the ellipse are that,

$$\cos \theta = -\frac{p}{r} = -\frac{p'}{r'} = -\sqrt{\frac{pp'}{rr'}} = -\frac{b}{\sqrt{rr'}} = \frac{-b}{\sqrt{a^2 - e^2x^2}} = \frac{-ab^2}{\sqrt{b^4x^2 + a^4y^2}}.$$

And, since on the ellipse $b^2xx' + a^2yy' = 0$, the numerator in the first member of (5) has the value

$$\frac{b^4x'^2}{y^2} = \frac{a^4y'^2}{x^2}.$$

Hence (5) implies

$$(6_1) \quad x'^2 = \frac{a\alpha y^2}{b^2}, \quad (6_2) \quad y'^2 = \frac{b^2\alpha x^2}{a^3};$$

whence

$$v^2 = \frac{\alpha(a^4y^2 + b^4x^2)}{a^3b^2} = \frac{\alpha rr'}{a},$$

or

$$(7) \quad \alpha = \frac{av^2}{rr'},$$

which is a generalization of (4).

By differentiating (6₁) as to time, and setting in the value of y' from (6₂), we have,

$$x'' = \frac{1}{2a} \frac{d\alpha}{dx} (a^2 - x^2) - \frac{x\alpha}{a}.$$

But the x -component of acceleration is

$$x'' = \alpha \frac{ae - x}{a - ex}.$$

From the comparison of the last two equations, we have

$$\frac{d\alpha}{2\alpha} = \frac{e dx}{a - ex}.$$

Whence the Newtonian law:

$$(8) \quad \alpha = \frac{k^2}{(a - ex)^2} = \frac{k^2}{r^2}.$$

A comparison of (7) and (8) yields the following elegant form of the *vis viva* equation:

$$v^2 = \frac{k^2}{a} \frac{r'}{r}.$$

The above discussion of elliptical motion, with only the slightest modification is valid for hyperbolic and parabolic motion under a central force directed toward a focus. And the last equation is valid in all three cases without modification, r'/a having the limiting value 2 in the case of the parabola.


III. For the path of a projectile, we may take a parabola of higher order and let

$$f(x, y) = y - (a_0 + a_1x + a_2x^2 + \dots + a_nx^n) = y - \varphi(x).$$

The range is assumed so short that the vertical direction may be regarded as constant. Now α is the resultant of the vertical gravitational acceleration $-g(y)$, and a tangential acceleration α_t . Obviously $\alpha \cos \theta = -g \cos \psi$, where ψ is the angle between the normal and the vertical. Hence (3) for this example is, by reference to (1),

$$-\frac{\varphi''(x)x'^2}{\sqrt{f_x'^2 + f_y'^2}} = g \cos \psi = g \frac{dx}{\sqrt{dx^2 + dy^2}} = g \frac{f_y'}{\sqrt{f_x'^2 + f_y'^2}}.$$

or,



Therefore,

(9)

and,

$$x' = \frac{\sqrt{g}}{[-\varphi''(x)]^{1/2}}.$$

$$v = \frac{\sqrt{g}}{[-\varphi''(x)]^{1/2}} \sqrt{1 + \left(\frac{dy}{dx}\right)^2};$$

$$x'' = \frac{g}{2} \frac{\varphi'''(x)}{[\varphi''(x)]^2}.$$

But since the total horizontal acceleration is the horizontal component of α_t , it follows that

$$(10) \quad \alpha_t = \frac{g\varphi'''(x)}{2[\varphi''(x)]^2} \cdot \sqrt{1 + \left(\frac{dy}{dx}\right)^2};$$

or, by (9),

$$\alpha_t = v \frac{\sqrt{g}\varphi'''(x)}{2[-\varphi''(x)]^{3/2}}.$$

Now (9) and (10) can be written in the form $v = F_1(x)$ and $\alpha_t = F_2(x)$, where F_1 and F_2 are power series. The method of inversion of series gives

$$(11) \quad F_2^{-1}(\alpha_t) = F_1^{-1}(v).$$

This entire discussion of the motion of a projectile is valid if the projectile is a finite sphere and also if the medium is not homogeneous. But the relation of α_t to v depends on the character and condition of the medium. So, (11) has the nature of a law for the particular projectile if, and only if, the medium in which the flight occurred was equally resistant at all points in the observed path.